

A Composite Fermion Hofstadter Problem: Partially Polarized Density Wave States in the $\nu = 2/5$ FQHE

Ganpathy Murthy

Physics Department, Boston University, Boston MA 02215

and Department of Physics and Astronomy, Johns Hopkins University, Baltimore MD 21218

(February 1, 2008)

It is well known that the $\nu = 2/5$ state is unpolarized at zero Zeeman energy, while it is fully polarized at large Zeeman energies. A novel state with charge/spin density wave order for Composite Fermions is proposed to exist at intermediate values of the Zeeman coupling for $\nu = 2/5$. This state has half the maximum possible polarization, and can be extended to other incompressible fractions. A Hartree-Fock calculation based on the new approach for all fractional quantum Hall states developed by R. Shankar and the author is used to demonstrate the stability of this state to single-particle excitations, and compute gaps. We compare our results with a very recent experiment which shows direct evidence for the existence of such a state, and also with more indirect evidence from past experiments.

73.50.Jt, 05.30.-d, 74.20.-z

The fractional quantum Hall (FQH) effect¹ has introduced us to new, highly correlated, incompressible states² of electrons in high magnetic fields. A unified understanding of all fractions $\nu = p/(2sp+1)$ was achieved by the Composite Fermion picture³, in which the electrons are dressed by $2s$ units of statistical flux to form Composite Fermions (CFs). At a mean field level, the CFs see a reduced field $B^* = B/(2sp+1)$, in which they fill p CF-Landau levels (CF-LLs), and exhibit the integer quantum Hall effect.

Due to the small g factor of electrons in GaAs, spins may not be fully polarized in FQH states^{4,5}. Transitions between singlet, partially polarized, and fully polarized states (based on gap measurements) have been observed for a number of fillings⁶⁻⁹, which can be understood in terms of CF's with a spin^{3,9,10}. The transitions happen when an unoccupied CF-LL of one spin crosses the occupied CF-LL of the opposite spin. Until recently, detailed measurements of the spin polarization in a FQH state were restricted to $1/3$ ¹¹.

Very recently, Kukushkin *et al* have reported direct measurements of the total spin polarization for a number of FQH states¹². In addition to strong transitions associated with CF-LL crossings, with plateaus of the proper polarizations predicted by CF theory¹⁰, they observe weaker plateaus which cannot be explained within the current framework⁹. At $\nu = 2/5$ they observe plateaus corresponding to half the maximal polarization at intermediate values of the Zeeman coupling¹². These odd features make their appearance in all the incompressible fractions studied.

I propose a partially polarized density wave (PPDW) state of CFs which is consistent with the experimental observations. The idea is this: At $\nu = 2/5$ the Zeeman energy $E_Z = g\mu B_{tot}$ favors polarization, while the electron-electron interaction favors a singlet state^{5,10}. By forming a periodic structure with two quanta of *effective* flux per unit cell, each CF-LL gets split into two sub-

bands. The $n = 0$ $\downarrow$ -spin CF-LL and the $n = 1$ $\uparrow$ -spin CF-LLs can both be half-occupied at intermediate E_Z , resulting in half the maximal polarization. The $n = 0$ $\uparrow$ -spin band is always fully occupied. When the chemical potential is in the subband gap the polarization is independent of E_Z , leading to a polarization plateau. Such a state can also have a quantized Hall conductance (by forming a "Hall crystal"¹³).

Charge density wave (CDW) states of electrons in high magnetic fields have been explored in Hartree-Fock (HF) theory¹⁴⁻¹⁷, in which the interacting problem is converted into a single-electron problem in a periodic potential, *i.e.*, the Hofstadter problem. In the strongly correlated lowest LL (LLL) CDW states of electrons are not attained until very low densities ($\nu < 1/5$)¹⁸.

In order to substantiate the idea proposed above, a computational scheme which gives reliable results for arbitrary FQH states is needed. Recently R. Shankar and the present author have developed just such an approach¹⁹, based on previous functional descriptions²⁰. Our central result is a formula for the LLL-projected electronic charge density at small q :

$$\rho_e(q) = \frac{\sum_j e^{-iqx_j}}{2ps+1} - il^2 \left(\sum_j (q \times \Pi_j) e^{-iqx_j} \right) \quad (1)$$

where $\vec{x}_j$ is a CF coordinate, $l = 1/\sqrt{eB}$ is the magnetic length, and $\vec{\Pi}_j = \vec{P}_j + e\vec{A}^*(r_j)$ is the velocity operator of the CFs. The low-energy Hamiltonian is $H = \frac{1}{2} \int \frac{d^2q}{(2\pi)^2} v(q) \bar{\rho}(-q) \bar{\rho}(q)$ where $v(q)$ is the electron-electron interaction. To include the effects of finite sample thickness, and to stay within the limitations of our small- q approach, we work with a modified Coulomb interaction of the form $v(q) = e^{-\lambda q} 2\pi e^2/q$, where the length λ is connected to the thickness. In earlier papers we presented HF calculations of gaps for a few fractions²¹, tested certain scaling relations that arise naturally in our theory against CF-wavefunction

results²², and computed magnetoexciton dispersions²³ in the TDHF approximation. The results show that HF in terms of CF variables gives a reasonably good account of physical properties.

The reason HF is good here is that in our formulation we are working directly with the quasiparticles of the system, the CFs. These are quasiparticles in the same sense as in Landau's Fermi liquid theory: they may interact strongly, but the interaction only scatters them weakly out of their states at low energies, making a description in terms of single-particle CF states applicable. Thus we expect the HF calculation to be described below to be good to the same level of accuracy (about 20%).

We start with the wavefunctions for the n^{th} CF-LL

$$\phi_{n,X}(\vec{r}) = \frac{e^{\frac{iyX}{l^{*2}} - \frac{(x-X)^2}{2l^{*2}}}}{\sqrt{l^*L\sqrt{\pi}2^n n!}} H_n\left(\frac{x-X}{l^*}\right) \quad (2)$$

Here $\vec{r}$ is a CF coordinate, L is the linear size of the system (area = L^2), and $l^* = l\sqrt{2ps+1}$ is the magnetic length in the effective field. X is an integer multiple of $\frac{2\pi l^{*2}}{L}$, and H_n is a Hermite polynomial. The degeneracy of each CF-LL is $\frac{L^2}{2\pi l^{*2}}$, which is $1/(2ps+1)$ times the degeneracy of the electronic LLL. Using standard manipulations we can express the momentum space electron density operator as

$$\hat{\rho}(\vec{q}) = \sum_{n_1, n_2, X} e^{-iq_x X} a_{n_1, X-}^\dagger a_{n_2, X+} \rho_{n_1 n_2}(\vec{q}) \quad (3)$$

where $X_\pm = X \pm \frac{q_y l^{*2}}{2}$, $a_{n, X}$ destroys a CF in the state $\phi_{n, X}$, and the final matrix element is given by

$$\rho_{n_1 n_2}(\vec{q}) = \frac{1}{2^{p+1}} e^{i\theta_q(n_1 - n_2) - i\frac{\pi}{2}(n_g - n_l)} \sqrt{\frac{n_l!}{n_g!}} \left(\frac{q l^*}{\sqrt{2}}\right)^{n_g - n_l} \times e^{-\frac{q^2 l^{*2}}{4}} (n_g L_{n_l-1}^{n_g - n_l} + 2L_{n_l}^{n_g - n_l} - (n_l + 1)L_{n_l+1}^{n_g - n_l}) \quad (4)$$

where θ_q is the angle $\vec{q}$ makes with the x -axis, n_g and n_l are the greater and lesser of n_1, n_2 , and the L_n^k are the Laguerre polynomials whose argument is $q^2 l^{*2}/2$.

The Hamiltonian can now be written as

$$H = \frac{1}{2L^2} \sum_{\vec{q}\sigma\sigma', \{n_i\}, \{X_i\}} v(\vec{q}) e^{-iq_x(X_1 - X_2)} \rho_{n_1 n_2}(\vec{q}) \rho_{n_3 n_4}(-\vec{q}) a_{\sigma, n_1, X_1-}^\dagger a_{\sigma, n_2, X_1+} a_{\sigma', n_3, X_2+}^\dagger a_{\sigma', n_4, X_2-} \quad (5)$$

where σ, σ' represent the spin labels²⁴. The electronic hamiltonian in the LLL is very similar, except that it must be normal-ordered to prevent self-interactions. Eq.(5) above is the correct form for the CF hamiltonian¹⁹, and the energy coming from normal-ordering represents the Hartree interaction of an electron with its own correlation hole.

We allow for a density wave by making the ansatz¹⁴

$$\langle a_{\sigma, n_1, X - \frac{Q_y l^{*2}}{2}}^\dagger a_{\sigma, n_2, X + \frac{Q_y l^{*2}}{2}} \rangle = e^{iQ_x X} \Delta_{\sigma, n_1 n_2}(\vec{Q}) \quad (6)$$

In principle, one should allow for nonzero Δ for all values of the subscripts, for all $\vec{Q}$'s consistent with the assumed lattice structure, and repeat for all possible lattice structures. We will carry out a restricted HF on a square lattice where only $\Delta_{\sigma, nn}$ for $n = 0, 1$ are assumed nonzero, thus ignoring CF-LL mixing. We consider only the smallest reciprocal vectors $\vec{Q}_1 = \pm Q_0 \hat{e}_x$, $\pm Q_0 \hat{e}_y$ with $\Delta = \Delta_1$, and the next smallest vectors $\vec{Q}_2 = Q_0(\pm \hat{e}_x \pm \hat{e}_y)$, $Q_0(\pm \hat{e}_x \mp \hat{e}_y)$ with $\Delta = \Delta_2$. It is essential to consider $\vec{Q}_2$ for the square lattice, because without them there is no gap between the two subbands, and no partially polarized plateau will be obtained.

I believe no physics is being lost by these restrictions. For half-filling, the square lattice has a lower HF energy than the triangular one¹⁴, and only the above two kinds of reciprocal vectors need be kept for an accurate treatment¹⁴. Also, including (CF-)LL mixing does not change the physics¹⁷.

Since the $n = 0$ $\uparrow$ -spin CF-LL is always fully occupied, $\Delta_{\uparrow, 00} = 0$ for all $\vec{Q}$. There are thus 4 variables $\Delta_{1\uparrow}, \Delta_{2\uparrow}, \Delta_{1\downarrow}, \Delta_{2\downarrow}$. The full HF hamiltonian can be written as

$$H_{HF} = \sum_{\sigma, n, X} \epsilon_{\sigma, n}^0 a_{\sigma, n, X}^\dagger a_{\sigma, n, X} + \sum_{\sigma, X, \vec{Q}_i} V_{i\sigma} a_{\sigma, i, X+}^\dagger a_{\sigma, i, X-} e^{iQ_x X} \quad (7)$$

where, in units of $e^2/\epsilon l^*$

$$\epsilon_{\sigma, n}^0 = \int_0^\infty d\tilde{q} \frac{e^{-\lambda^* \tilde{q}}}{2} \sum_{n'=0}^{n_{max}} (1 - 2N_F(\sigma, n')) |\rho_{nn'}(\vec{q})|^2 \quad (8)$$

$$V_{i, \sigma} = \Delta_{i, -\sigma} \frac{e^{-\lambda^* Q_i} \rho_{00}(Q_i) \rho_{11}(Q_i)}{Q_i l^*} + \Delta_{i, \sigma} \left(\frac{e^{-\lambda^* Q_i} \rho_{mm}(Q_i)^2}{Q_i l^*} - \int_0^\infty d\tilde{q} e^{-\lambda^* \tilde{q}} J_0(\tilde{q} Q_i l^*) \rho_{mm}(\tilde{q})^2 \right) \quad (9)$$

where $\tilde{q} = q l^*$, $\lambda^* = \lambda/l^*$, $m = 1$ for the $\uparrow$ -spin, and $m = 0$ for the $\downarrow$ -spin, and $N_F(\sigma, n')$ represents the Fermi occupation of the spin- σ n' CF-LL. In our problem of interest, $N_F(\uparrow, 0) = 1$, $N_F(\uparrow, 1) = N_F(\downarrow, 0) = 1/2$.

Now, one diagonalizes the HF-hamiltonian by a sequence of unitary transformations¹⁶ for the case of rational effective flux through a unit cell. For the square lattice this is $\frac{Q_0^2 l^{*2}}{2\pi} = \frac{M}{N}$. Each CF-LL splits into N subbands with equal degeneracies. Since we are interested in the half-filled CF-LL case, we minimally choose $M = 1$, $N = 2$, with two quanta of effective flux through each unit cell. For this simple case one can write down the energies analytically. They turn out to be $\epsilon_\sigma = \epsilon_\sigma^0 \pm \omega_\sigma$ where

$$\omega_\sigma = 2 \left(V_{1\sigma}^2 (\cos^2(Q_0 X) + \cos^2(Q_0 Y)) + 4V_{2\sigma}^2 \sin^2(Q_0 X) \sin^2(Q_0 Y) \right)^{\frac{1}{2}} \quad (10)$$

where $\sigma = \uparrow$ implies the $n = 1$ CF-LL, and $\sigma = \downarrow$ the $n = 0$ CF-LL. X, Y are integer multiples of $\frac{2\pi l^{*2}}{L}$ which range from 0 to $\sqrt{\pi}$. Finally we impose the self-consistency conditions

$$\begin{aligned}\Delta_{1\sigma} &= -\frac{V_{1\sigma}}{\pi Q_0} \int_0^\pi d\tilde{X} d\tilde{Y} \frac{\cos^2(\tilde{X})}{\omega_s} \\ \Delta_{2\sigma} &= -\frac{2V_{2\sigma}}{\pi Q_0^2} \int_0^\pi d\tilde{X} d\tilde{Y} \frac{\sin^2(\tilde{X}) \sin^2(\tilde{Y})}{\omega_s}\end{aligned}\quad (11)$$

The gap between the two subbands is $8|V_{2\sigma}|$. Let us denote the top of the occupied subband as the highest occupied molecular orbital (HOMO), and the bottom of the unoccupied subband as the lowest unoccupied molecular orbital (LUMO). In the absence of Zeeman energy I find that $E_{HOMO}^\uparrow > E_{LUMO}^\downarrow$, which is inconsistent; eq.(11) is predicated on the chemical potential lying in both the subband gaps, which must therefore intersect. As the Zeeman energy increases the entire $n = 0$ $\downarrow$ -spin structure moves up, while the $n = 1$ $\uparrow$ -spin structure moves down. There is a lower critical Zeeman energy $E_Z^<$ for which the two subband gaps intersect. As the Zeeman energy increases further eventually we will have an upper critical Zeeman energy $E_Z^>$ where $E_{LUMO}^\uparrow = E_{HOMO}^\downarrow$. For $E_Z > E_Z^>$ the proposed state is again inconsistent. The PPDW state exists and is stable (to single-particle excitations) for $E_Z^< < E_Z < E_Z^>$.

Now for the numerical results. I have found self-consistent solutions for $\lambda = l, 1.5l$, since these values are likely to bracket the thickness of physical samples²⁵. I find that the PPDW state is locally stable for $0.006 < E_Z < 0.012$ at $\lambda = l$, while for $\lambda = 1.5l$ the regime of stability is $0.002 < E_Z < .01$. Here and below E_Z and all gaps are in units of $e^2/\epsilon l$.

The strongest experimental evidence for PPDW states comes from the total polarization measurements of Kukushkin *et al*¹². Examining their curve for $\nu = 2/5$ we see that the partially polarized state with a constant polarization of half the maximum value is seen for $0.009 < E_Z < 0.011$. This is fully consistent with our results for $\lambda = l$, since all we have ascertained is the stability of the PPDW state to single-particle excitations, and ignored other instabilities.

Evidence for PPDW states is seen in their data for other incompressible fractions as well¹². For example, their curve for $\nu = 2/3$ shows a small region of polarization 1/2, which is impossible to explain using translationally invariant states of CFs. Similar regions are seen in the 4/7, 3/7, and 4/9 data. This leads us to believe that the PPDW state is a generic feature of the incompressible fractions where the ground state is not fully polarized at $E_Z = 0$. While the polarization measurement is unambiguous, it would be nice to obtain direct evidence of the inhomogeneity, perhaps by surface acoustic wave measurements²⁶. The period for the square lattice is $2l^*\sqrt{\pi} = 7.93l$.

There is also suggestive evidence from earlier experiments. For 4/3, 8/5, and 7/5, Du *et al*⁹ find that while the gap seems to go to zero at the transition, it does not come back up to the expected value, but remains (for the latter two fractions) an order of magnitude below. Theoretically, gaps in the PPDW state are subband gaps, and are naturally small. For example, at $\lambda = l$ in our approach, the singlet state has a gap of 0.045, the fully

polarized state has a gap 0.032, while the partially polarized state gap (which changes with E_Z) has a maximum value of 0.0029, an order of magnitude below the usual FQHE gaps. An earlier study⁶ found a “gapless” region at intermediate E_Z for 4/3 beyond which the gap recovered. Presumably the gap was below the resolution in that experiment, since a later experiment⁸ found quantization of σ_{xy} in the “gapless” region. Du *et al* also find maxima in the longitudinal resistance not only at the CF-LL level crossings (the strong maxima), but also at other values of E_Z (which are weaker maxima), which they are unable to understand⁹. These weaker maxima may be caused by transitions to PPDW states.

Finally, there is some evidence from exact diagonalizations of small systems²⁷ that the 3/5 and 4/9 states have a “gapless” regime at intermediate E_Z . The period of the PPDW state is too large to show up in these studies.

Let us turn to some of the caveats concerning our calculation, where many effects are ignored, including CF-LL mixing and larger $\vec{Q}$'s. We have concentrated on 2 units of effective flux per unit cell. This is symmetric and natural in this problem since we need the $\uparrow$ -spin CF-LL with filling ν_1 , and the $\downarrow$ -spin CF-LL with filling $1 - \nu_1$ to have large gaps. This is also what is seen in the data¹². However, other configurations are possible. We have treated the constraint by cutting off the number of CF-LLs ($n_{max} = 4$) as in our previous calculations^{21,23}. As the fractions approach a compressible state, a better treatment of the constraint will become necessary²⁸. We have used the small- q version of the density, whereas the actual value is $Q_0 \approx l^{-1}$. This can be remedied by using a recent extension of our formulation²⁹ to all q . If a charge density inhomogeneity exists, the CF's will feel a periodic Chern-Simons field due to the charge texture. The present treatment produces weak charge modulation ($\rho(Q_0)/\rho(0) \approx 0.01$), and we have ignored this effect³⁰. We have only considered the stability to single particle excitations, while there may be other instabilities as well. One omission is significant: We have not compared ground state energies of the various states. The reason is that the ground state structure factor $S(k)$ in our approach is not correct (at small k)³¹. Our approach is an expansion in gradient interactions¹⁹. Due to the peculiar properties of the electronic density matrix elements, the order we have kept suffices for reasonably accurate calculations of gaps, but one needs to go to higher order to get $S(k)$ accurately¹⁹.

Let us now explore some potential generalizations of our results. An obvious extension is to the $\nu = 2$ state. My preliminary results indicate that the PPDW state should be locally stable here as well³⁵. To access this region experimentally would require huge in-plane fields, or very low densities. A loose analogy can also be made between the 2/5 two CF-LL problem and the bilayer quantum Hall problem³², with the CF-LL index playing the role of the layer index. The two fully filled CF-LLs in $\nu = 2/5$ correspond to the $\nu = 2$ bilayer problem, for which a translationally invariant canted antiferromagnetic state has been proposed³³ at intermediate E_Z . A general 3-parameter wavefunction has been written for

such states³⁴, which should compete with the PPDW state. (For $\nu = 2/5$ in our formalism, the analog of this state³⁴ seems to have no HF solutions other than the singlet and fully polarized states³⁵: In any case, the canted states should show a smooth variation of the total polarization with E_Z , not the plateaus seen in the data¹²). If we push the analogy further, the PPDW state should occur in the bilayer $\nu = 2$ case for the two layers having unequal fillings $1/2$ and $3/2$. These generalizations will be explored elsewhere³⁵.

In summary, I have presented theoretical arguments in support of a partially polarized density wave state of Composite Fermions at $\nu = 2/5$ (and other incompressible fractions), which can be accessed by tuning the Zeeman energy. I have also argued that such states have already been seen in a recent experiment¹², and indicated suggestive features of older experiments which hint at the existence of these states. It would be fascinating to explore the connections to other inhomogeneous states proposed in the quantum Hall regime^{13,36,37}.

It is a pleasure to thank H.A.Fertig, J.K.Jain, and Z.Tesanovic for illuminating conversations.

-
- ¹ D.Tsui, H.Stromer and A.Gossard, Phys. Rev. Lett. **48**, 1599, (1982).
 - ² R.B.Laughlin, Phys. Rev. Lett. **50**, 1395, (1983)
 - ³ J.K.Jain, Phys. Rev. Lett. **63**, 199, (1989); Phys. Rev. **B 41**, 7653 (1990); Science **266**, 1199 (1994); J.K.Jain and R.K.Kamilla, in "Composite Fermions", Olle Heinonen, Editor (World Scietific, 1998).
 - ⁴ B.I.Halperin, Helv. Phys. Acta **556**, 75 (1983).
 - ⁵ T.Chakraborty and F.C.Zhang, Phys. Rev. **B29**, 7032; F.C.Zhang and T.Chakraborty, *ibid* **30** 7320; M.Rasolt, F.Perrot, and A.H.MacDonald, Phys. Rev. Lett.**55**, 433 (1985).
 - ⁶ R.G.Clark, S.R.Haynes, A.M.Suckling, J.R.Mallett, P.A.Wright, J.J.Harris, and C.T.Foxon, Phys. Rev. Lett.**62**, 1536 (1989).
 - ⁷ J.P.Eisenstein, H.L.Stormer, L.Pfeiffer, and K.W.West, Phys. Rev. Lett.**62**, 1540 (1989); J.E.Furneaux, D.A.Syphers, and A.G.Swanson, Phys. Rev. Lett.**63**, 1098 (1989).
 - ⁸ A.Buckthought, R.Boulet, A.Sachrajda, Z.Wasilewski, P.Zawadzski, and F.Guillon, Sol. St. Comm. **78**, 191 (1991).
 - ⁹ R.R.Du, A.S.Yeh, H.L.Stormer, D.C.Tsui, L.N.Pfeiffer, and K.W.West, Phys. Rev. Lett.**75**, 3926 (1995); Phys. Rev. **B55**, R7351 (1997).
 - ¹⁰ X.G.Wu, G.Dev, and J.K.Jain, Phys. Rev. Lett.**71**, 153 (1993); K.Park and J.K.Jain, Phys. Rev. Lett.**80**, 4237 (1998).
 - ¹¹ P.Khandelwal, N.N.Kuzma, S.E.Barrett, L.N.Pfeiffer, and K.W.West, Phys. Rev. Lett.**81**, 673 (1998).
 - ¹² I.V.Kukushkin, K.v Klitzing, and K.Eberl, Phys. Rev. Lett.**82**, 3665 (1999).
 - ¹³ B.I.Halperin, Z.Tesanovic, and F.Axel, Phys. Rev. Lett.**57**, 922 (1986); Z.Tesanovic, F.Axel, and B.I.Halperin, Phys. Rev. **B39**, 8525 (1989); A.Kol and N.Read, Phys. Rev. **B48**, 8890 (1993).
 - ¹⁴ D.Yoshioka and H.Fukuyama, Jour. Phys. Soc. Japan **47**, 394 (1979).
 - ¹⁵ H.Fukuyama, P.M.Platzman, and P.W.Anderson, Phys. Rev. **B19**, 5211 (1979); H.Fukuyama and P.M.Platzman, Phys. Rev. **B25**, 2934 (1982).
 - ¹⁶ D.Yoshioka and P.A.Lee, Phys. Rev. **B27**, 4986 (1983).
 - ¹⁷ A.H.MacDonald, Phys. Rev. **B30**, 4392 (1984).
 - ¹⁸ For a review of the Wigner Crystal, see H.A.Fertig, in "Perspectives in Quantum Hall Effects", S.Das Sarma and A.Pinczuk editors (Wiley, 1997).
 - ¹⁹ R.Shankar and G.Murthy, Phys. Rev. Lett. **79**, 4437, (1997); G.Murthy and R.Shankar, in "Composite Fermions", Olle Heinonen, Editor, (World Scietific, 1998).
 - ²⁰ S.M.Girvin, in "The Quantum Hall Effect", Edited by R.E.Prange and S.M. Girvin, (Springer-Verlag, 1990); S.M.Girvin and A.H.MacDonald, Phys. Rev. Lett. **58**, 1252 (1987); S.-C.Zhang, H.Hansson and S.A.Kivelson, Phys. Rev. Lett. **62**, 82, (1989); N.Read, Phys. Rev. Lett., **62**, 86 (1989); A.Lopez and E.Fradkin, Phys. Rev. B **44**, 5246 (1991), *ibid* **47**, 7080, (1993), Phys. Rev. Lett. **69**, 2126 (1992); V.Kalmeyer and S.-C.Zhang, Phys. Rev. B **46**, 9889 (1992); B.I.Halperin, P.A.Lee and N.Read, Phys. Rev. B **47**, 7312 (1993).
 - ²¹ G.Murthy and R.Shankar, Phys. Rev. **B59**, 12260 (1999).
 - ²² G.Murthy, K.Park, R.Shankar, and J.K.Jain, Phys. Rev. **B58**, 15363 (1998).
 - ²³ G.Murthy, cond-mat/9903187.
 - ²⁴ We are implicitly attaching flux independent of spin²³.
 - ²⁵ R.H.Morf, cond-mat/9812181.
 - ²⁶ See, for example, R.L.Willet, K.W.West, and L.N.Pfeiffer, Phys. Rev. Lett.**78**, 4478 (1997).
 - ²⁷ T.Chakraborty and P.Pietilainen, Phys. Rev. **B41**, 10862 (1990).
 - ²⁸ B.I.Halperin and A.Stern, Phys. Rev. Lett.**80**, 5457 (1998); G.Murthy and R.Shankar, *ibid* 5458 (1998); D.H.Lee, Phys. Rev. Lett.**80**, 4745 (1998); N.Read, Phys. Rev. **B58**, 16262 (1998); A.Stern, B.I.Halperin, F.von Oppen, and S.H.Simon, cond-mat/9812135.
 - ²⁹ R.Shankar, cond-mat/9903064.
 - ³⁰ For the periodic field problem, see, D.Yoshioka and Y.Iye, Jour. Phys. Soc. Japan **56**, 448 (1987).
 - ³¹ For example, the small- k $1/3$ structure factor is a factor of 2 off the exact result of S.M.Girvin, A.H.MacDonald and P.Platzman, Phys. Rev. **B 33**, 2481 (1986).
 - ³² K.Yang, K.Moon, L.Zheng, A.H.MacDonald, S.M.Girvin, D.Yoshioka, and S.-C.Zhang, Phys. Rev. Lett.**72**, 732 (1994); For a review of the experiments, see J.P.Eisenstein in "Perspectives in Quantum Hall Effects", S.Das Sarma and A.Pinczuk editors (Wiley, 1997).
 - ³³ L.Zheng, R.J.Radtke, and S.Das Sarma, Phys. Rev. Lett.**78**, 2453 (1997).
 - ³⁴ A.H.MacDonald, R.Rajaraman, and T.Jungwirth, cond-mat/9903318.
 - ³⁵ G.Murthy, in preparation.
 - ³⁶ L.Brey, H.A.Fertig, R.Cote, and A.H.MacDonald, Phys. Rev. Lett.**75**, 2562 (1995).
 - ³⁷ A.A.Koulakov, M.M.Folger, and B.I.Shklovskii, Phys. Rev. Lett.**76**, 499 (1996).